

Références Kits Kinases Assay - Advion Interchim Scientific

Une fois votre référence choisie, rendez-vous sur notre site www.interchim.com pour commander aisément vos besoins.

Référence	Description	Cond.	Descriptif
500240	BTK (Phospho-Tyr223) TR-FRET Assay Kit	480 Well	This BTK (Phospho-Tyr223) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This BTK (Phospho-Tyr223) TR-FRET Assay Kit is robust, with a Z' value of 0.74, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500240	BTK (Phospho-Tyr223) TR-FRET Assay Kit	96 Well	This BTK (Phospho-Tyr223) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This BTK (Phospho-Tyr223) TR-FRET Assay Kit is robust, with a Z' value of 0.74, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500241	BTK (Total) TR-FRET Assay Kit	480 Well	This BTK (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This BTK (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500241	BTK (Total) TR-FRET Assay Kit	96 Well	This BTK (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This BTK (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500242	BTK (Total) and BTK (Phospho-Tyr223) TR-FRET Assay Kit	480 Well	This BTK (Total) and BTK (Phospho-Tyr223) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total and phosphorylated protein levels in cells. This BTK (Total) and BTK (Phospho-Tyr223) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing 96 total and 384 phosphorylated protein wells. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.

500251	GSK3 β (Phospho-Ser9) TR-FRET Assay Kit	480 Well	This GSK3 β (Phospho-Ser9) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This GSK3 β (Phospho-Ser9) TR-FRET Assay Kit is robust, with a Z' value of 0.74, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500251	GSK3 β (Phospho-Ser9) TR-FRET Assay Kit	96 Well	This GSK3 β (Phospho-Ser9) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This GSK3 β (Phospho-Ser9) TR-FRET Assay Kit is robust, with a Z' value of 0.74, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500252	GSK3 β (Total) TR-FRET Assay Kit	480 Well	This GSK3 β (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This GSK3 β (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500252	GSK3 β (Total) TR-FRET Assay Kit	96 Well	This GSK3 β (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This GSK3 β (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500253	GSK3 β (Total) and (Phospho-Ser9) TR-FRET Assay	480 Well	This GSK3 β (Total) and GSK3 β (Phospho-Ser9) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total and phosphorylated protein levels in cells. This GSK3 β (Total) and GSK3 β (Phospho-Ser9) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing 96 total and 384 phosphorylated protein wells. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.

500254	JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit	480 Well	This JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit is robust, with a Z' value of 0.82, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500254	JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit	96 Well	This JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit is robust, with a Z' value of 0.82, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500255	JNK1/2/3 (Total) TR-FRET Assay Kit	480 Well	This JNK1/2/3 (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This JNK1/2/3 (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500255	JNK1/2/3 (Total) TR-FRET Assay Kit	96 Well	This JNK1/2/3 (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This JNK1/2/3 (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500256	JNK1/2/3 (Total) and JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit	480 Well	This JNK1/2/3 (Total) and JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total and phosphorylated protein levels in cells. This JNK1/2/3 (Total) and JNK1/2/3 (Phospho-Thr183/Tyr185) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing 96 total and 384 phosphorylated protein wells. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.

500257	LCK (Phospho-Tyr394) TR-FRET Assay Kit	480 Well	This LCK (Phospho-Tyr394) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This LCK (Phospho-Tyr394) TR-FRET Assay Kit is robust, with a Z' value of 0.72, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500257	LCK (Phospho-Tyr394) TR-FRET Assay Kit	96 Well	This LCK (Phospho-Tyr394) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This LCK (Phospho-Tyr394) TR-FRET Assay Kit is robust, with a Z' value of 0.72, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500258	MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit	480 Well	This MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit is robust, with a Z' value of 0.76, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500258	MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit	96 Well	This MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit is robust, with a Z' value of 0.76, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500259	MEK1 (Total) TR-FRET Assay Kit	480 Well	This MEK1 (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This MEK1 (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.

500259	MEK1 (Total) TR-FRET Assay Kit	96 Well	This MEK1 (Total) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total protein levels in cells. This MEK1 (Total) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 total protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500261	MEK1 (Total) and MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit	480 Well	This MEK1 (Total) and MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of total and phosphorylated protein levels in cells. This MEK1 (Total) and MEK1 (Phospho-Ser218/222) TR-FRET Assay Kit is suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing 96 total and 384 phosphorylated protein wells. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500268	p38α MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit	480 Well	This p38α MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This p38α MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit is robust, with a Z' value of 0.71, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500268	p38α MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit	96 Well	This p38α MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This p38α MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit is robust, with a Z' value of 0.71, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500269	p38αβγ MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit	480 Well	This p38αβγ MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This p38αβγ MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit is robust, with a Z' value of 0.73, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.

500269	p38 α β MAPK (Phospho-Thr180/Tyr182) TR-FRET	96 Well	This p38 α β MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This p38 α β MAPK (Phospho-Thr180/Tyr182) TR-FRET Assay Kit is robust, with a Z' value of 0.73, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500284	Src (Phospho-Tyr419) TR-FRET Assay Kit	480 Well	This Src (Phospho-Tyr419) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This Src (Phospho-Tyr419) TR-FRET Assay Kit is robust, with a Z' value of 0.74, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500284	Src (Phospho-Tyr419) TR-FRET Assay Kit	96 Well	This Src (Phospho-Tyr419) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This Src (Phospho-Tyr419) TR-FRET Assay Kit is robust, with a Z' value of 0.74, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500289	TBK1/NAK (Phospho-Ser172) TR-FRET Assay Kit	480 Well	This TBK1/NAK (Phospho-Ser172) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This TBK1/NAK (Phospho-Ser172) TR-FRET Assay Kit is robust, with a Z' value of 0.70, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
500289	TBK1/NAK (Phospho-Ser172) TR-FRET Assay Kit	96 Well	This TBK1/NAK (Phospho-Ser172) TR-FRET Assay Kit uses a homogeneous TR-FRET assay method amenable to rapid measurement of phosphorylated protein levels in cells. This TBK1/NAK (Phospho-Ser172) TR-FRET Assay Kit is robust, with a Z' value of 0.70, and suitable for screening a large number of samples. The signal is stable at room temperature for at least 24 hours, affording flexibility in read times. The amount of reagents provided is sufficient for testing either 96 or 480 phosphorylated protein wells, depending on the size of the kit. THUNDER™ TR-FRET Cell Signaling Assay Kits are based on Bioauxilium's enhanced proprietary time-resolved Förster resonance energy transfer (TR-FRET) technology.
701741	SPHK1 Assay Buffer (1X)	14 mL	SPHK1 Assay Buffer (1X) has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 1 Inhibitor Screening Assay Kit (Item No. 701740).

701742	SPHK1 Enzyme (human, recombinant)	50 µL	SPHK1 Enzyme (human, recombinant) has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 1 Inhibitor Screening Assay Kit (Item No. 701740).
701743	SPHK Substrate	1 ea	SPHK Substrate has been formulated and tested to work with Cayman's Assay Kits. Detailed instructions for its use are contained in the respective Cayman Assay Kit booklet.
701744	SPHK1 ATP	5 mL	SPHK1 ATP has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 1 Inhibitor Screening Assay Kit (Item No. 701740).
701745	SPHK1 Inhibitor (PF-543)	100 µL	SPHK1 Inhibitor (PF-543) has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 1 Inhibitor Screening Assay Kit (Item No. 701740).
701871	SPHK2 Enzyme (human, recombinant)	50 µL	SPHK2 Enzyme (human, recombinant) has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 2 Inhibitor Screening Assay Kit (Item No. 701870).
701872	SPHK1/2 Inhibitor (SLC5111312)	1 ea	SPHK1/2 Inhibitor (SLC5111312) has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 2 Inhibitor Screening Assay Kit (Item No. 701870).
701873	SPHK2 Assay Buffer (1X)	14 mL	SPHK2 Assay Buffer (1X) has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 2 Inhibitor Screening Assay Kit (Item No. 701870).
701874	SPHK2 ATP	5 mL	SPHK2 ATP has been tested and formulated to work exclusively with Cayman's Sphingosine Kinase 2 Inhibitor Screening Assay Kit (Item No. 701870).