

Références Proteines Kinases - Advion Interchim Scientific

Une fois votre référence choisie, rendez-vous sur notre site www.interchim.com pour commander aisément vos besoins.

| Référence | Description | Cond. | Purité | Descriptif |
|-----------|---|--------|----------------------------|--|
| 10011209 | Akt1 (human, recombinant) | 1 ea | ≥95% estimated by SDS-PAGE | Source: Active recombinant N-terminal His-tagged Akt1 expressed in insect cells • Amino acids: 1-480 • MW: 60.3 kDa |
| 10011670 | Soluble Epoxide Hydrolase (mouse recombinant) | 50 µg | ≥95% estimated by SDS-PAGE | Source: Active mouse recombinant N-terminal His-tagged protein purified from Sf21 cells • Mr: 64 kDa • sEH catalyzes the hydrolysis of exogenous and endogenous epoxides to vicinal diols. Endogenous substrates for sEH include epoxyeicosatrienoic acids (EETs) which are known for their vasodilatory effects as well as for their anti-inflammatory actions. |
| 10348 | Sphingosine Kinase 1 (human, recombinant) | 25 µg | ≥80% estimated by SDS-PAGE | Source: Human recombinant N-terminal His-tagged protein from Sf9 cells • SPHK1 catalyzes the production of sphingosine-1-phosphate, a lipid mediator with broad spectrum of biological activities including cell proliferation, survival, migration, cytoskeletal organization, and morphogenesis. |
| 22817 | TBK1 (human, recombinant) | 50 µg | ≥50% estimated by SDS-PAGE | Source: Active recombinant N-terminal His-tagged protein expressed in insect cells • Amino acids: AA: 1-729 (full length) • MW: 84 kDa |
| 30191 | Sphingosine Kinase 2 (human, recombinant) | 25 µg | ≥75% estimated by SDS-PAGE | Source: Active recombinant N-terminal His-tagged protein expressed in insect cells • Amino acids: 2-654 (full length) • MW: 72.9 kDa |
| 31843 | PDGFRα/CD140a (human, recombinant) | 1 mg | ≥90% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged PDGFRA expressed in HEK293 cells • Amino acids: 24-524 • MW: 82.9 kDa |
| 31843 | PDGFRα/CD140a (human, recombinant) | 100 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged PDGFRA expressed in HEK293 cells • Amino acids: 24-524 • MW: 82.9 kDa |
| 31845 | VEGFR2 Extracellular Domain (human, recombinant; aa 20-327) | 200 µg | ≥95% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged VEGFR2 expressed in HEK293 Cells • Amino acids: 20-327 • MW: 61.5 kDa |
| 31846 | ALK4 Extracellular Domain (human, recombinant) | 1 mg | ≥92% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged ALK4 expressed in HEK293 cells • Amino acids: 24-126 • MW: 13 kDa |
| 31846 | ALK4 Extracellular Domain (human, recombinant) | 100 µg | ≥92% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged ALK4 expressed in HEK293 cells • Amino acids: 24-126 • MW: 13 kDa |
| 31847 | ALK5 Extracellular Domain (human, recombinant) | 100 µg | ≥85% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-His-tagged ALK5 expressed in HEK293 cells • Amino acids: 34-125 • MW: 38.1 kDa |
| 32007 | HER2/ERBB2 Extracellular Domain (human, recombinant) | 1 mg | ≥90% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged HER/ERBB2 expressed in HEK293 cells • Amino acids: 23-652 • MW: 96.1 kDa |
| 32007 | HER2/ERBB2 Extracellular Domain (human, recombinant) | 100 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged HER/ERBB2 expressed in HEK293 cells • Amino acids: 23-652 • MW: 96.1 kDa |
| 32026 | EGFR Extracellular Domain (human, recombinant) | 1 mg | ≥97% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged EGFR extracellular domain expressed in HEK293 cells • Amino acids: 25-645 • MW: 95 kDa |
| 32026 | EGFR Extracellular Domain (human, recombinant) | 100 µg | ≥97% estimated by SDS-PAGE | Source: Active recombinant C-terminal human IgG1 Fc-tagged EGFR extracellular domain expressed in HEK293 cells • Amino acids: 25-645 • MW: 95 kDa |
| 32557 | AMPK Complex (α1, β1, and γ1 subunits; human, recombinant) | 10 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged AMPK complex (α1, β1, and γ1 subunits) expressed in insect cells • Amino acids: 1-559, 1-270, and 1-331 for the α1, β1, and γ1 subunits, respectively • MW: ~146 kDa |
| 32563 | PKCα (human, recombinant) | 10 µg | ≥80% estimated by SDS-PAGE | Source: Active recombinant human N-terminal GST-tagged PKCα expressed in insect cells • Amino acids: 1-672 (full length) • MW: 102 kDa |

| | | | | |
|-------|---|--------|--------------------------------|--|
| 32564 | PKM2 (human, recombinant) | 20 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human N-terminal His-tagged PKM2 expressed in E. coli • Amino acids: 1-531 (full length) • MW: 58.9 kDa |
| 32565 | Src (human, recombinant) | 10 µg | ≥60% as determined by SDS-PAGE | Source: Active recombinant human N-terminal GST-tagged Src expressed in insect cells • Amino acids: 1-536 • MW: 86 kDa |
| 32757 | ROCK1 (human, recombinant) | 10 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant N-terminal His-tagged human ROCK1 expressed in insect cells • Amino acids: 6-415 • MW: 50.12 kDa |
| 32758 | ROCK2 (human, recombinant) | 10 µg | ≥95% | Source: Active recombinant human N-terminal His-tagged ROCK2 expressed in insect cells • Amino acids: 19-417 • MW: 48.3 kDa |
| 33731 | ALK7 Extracellular Domain (human, recombinant) | 200 µg | ≥95% estimated by SDS-PAGE | Source: Recombinant C-terminal human IgG1 Fc-tagged ALK7 expressed in HEK293 cells • Amino acids: 22-113 • MW: 36.6 kDa |
| 33735 | FGFR1 Extracellular Domain (human, recombinant) | 1 mg | ≥98% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged FGFR1 expressed in HEK293 cells • Amino acids: 22-285 • MW: 31 kDa |
| 33735 | FGFR1 Extracellular Domain (human, recombinant) | 100 µg | ≥98% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged FGFR1 expressed in HEK293 cells • Amino acids: 22-285 • MW: 31 kDa |
| 33736 | FLT3/CD135 Intracellular Domain (human, recombinant) | 10 µg | ≥80% as determined by SDS-PAGE | Source: Active recombinant human N-terminal His-tagged FLT3/CD135 expressed in insect cells • Amino acids: 564-993 • MW: 55 kDa |
| 33738 | GSK3β (human, recombinant) | 10 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human GSK-3β expressed in insect cells • Amino acids: 2-433 (full length) • MW: 75 kDa |
| 33739 | IKKβ (human, recombinant) | 10 µg | ≥80% estimated by SDS-PAGE | Source: Active recombinant human N-terminal GST-tagged IKKβ expressed in insect cells • Amino acids: 1-756 (full length) • MW: 105 kDa |
| 33741 | JAK1 (human, recombinant) | 10 µg | ≥63% estimated by SDS-PAGE | Source: Active recombinant human N-terminal GST-tagged JAK1 expressed in insect cells • Amino acids: 866-1,154 • MW: 60 kDa |
| 33742 | JAK2 (human, recombinant) | 20 µg | ≥71% estimated by SDS-PAGE | Source: Active recombinant human N-terminal His-tagged JAK2 expressed in insect cells • Amino acids: 808-1,132 • MW: 42.5 kDa |
| 33743 | MERTK Extracellular Domain (human, recombinant) | 10 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human N-terminal GST-tagged MERTK expressed in insect cells • Amino acids: 578-872 • MW: 60 kDa |
| 33745 | TYK2 (human, recombinant) | 10 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human N-terminal His-tagged TYK2 expressed in insect cells • Amino acids: 871-1,187 • MW: 38 kDa |
| 33746 | VEGFR3 Extracellular Domain (human, recombinant) | 1 mg | ≥97% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged VEGFR3 expressed in HEK293 cells • Amino acids: 25-776 • MW: 86 kDa |
| 33746 | VEGFR3 Extracellular Domain (human, recombinant) | 100 µg | ≥97% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged VEGFR3 expressed in HEK293 cells • Amino acids: 25-776 • MW: 86 kDa |
| 33986 | PDGFRβ/CD140b Extracellular Domain (human, recombinant) | 1 mg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged PDGFRβ expressed in HEK293 cells • Amino acids: 33-531 • MW: 57.6 kDa |
| 33986 | PDGFRβ/CD140b Extracellular Domain (human, recombinant) | 100 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human C-terminal His-tagged PDGFRβ expressed in HEK293 cells • Amino acids: 33-531 • MW: 57.6 kDa |
| 33987 | RET Extracellular Domain (human, recombinant) | 10 µg | ≥90% estimated by SDS-PAGE | Source: Active recombinant human N-terminal GST-tagged RET expressed in insect cells • Amino acids: 658-1114 • MW: 77.5 kDa |
| 33989 | PI3K p110γ (human, recombinant) | 20 µg | ≥80% estimated by SDS-PAGE | Source: Active recombinant human N-terminal His-tagged PI3K p110γ expressed in insect cells • Amino acids: 1-1,102 (full length) • MW: 130 kDa |
| 33990 | TAK1-TAB1 Fusion (human, recombinant) | 10 µg | ≥93% estimated by SDS-PAGE | Source: Active recombinant N-terminal GST-tagged human TAK1 fused to TAB1 via a DFGGGGG peptide linker expressed in insect cells • Amino acids: TAK1 (1-303) and TAB1 (437-504) • MW: 69 kDa |